

## Towards risk-adapted therapy for rhabdoid tumour subgroups

Atypical teratoid rhabdoid tumour is an aggressive paediatric embryonal brain tumour with a poor prognosis despite intensive multimodal therapy. Children older than 3 years with complete tumour resection and early irradiation might have better chances of survival.<sup>1</sup> The characteristics of atypical teratoid rhabdoid tumours are a biallelic inactivation of *SMARCB1* (*hSNF5/INI1/BAF47*), a core subunit of the highly-conserved SWI/SNF chromatin-remodelling complexes that have different roles in transcriptional and epigenetic regulation of cell signalling, growth, and differentiation.<sup>2</sup> First recognised in malignant rhabdoid tumours, inactivating mutations of SWI/SNF subunits have since been discovered in many cancers. Despite its overtly malignant phenotype, atypical teratoid rhabdoid tumour has a remarkably stable genome.<sup>3</sup> Thus attention has focused on epigenetic tumourigenic mechanisms underlying *SMARCB1* mutation, the only recurrent genetic alteration in atypical teratoid rhabdoid tumours.

Loss of *SMARCB1* leads to epigenetic alterations that drive key developmental and cancer-promoting pathways. In a normal state, *SMARCB1* interacts with epigenetic modifiers to suppress oncogenes and stimulate tumour suppressors (eg, recruitment of histone deacetylases to repress *cyclin D15* and evicts polycomb complex proteins to enhance transcription of *p15<sup>INK4b</sup>* or *p16<sup>INK4a</sup>*). *SMARCB1* also directly affects chromatin structure at gene promoters (eg, suppresses *GLI4* and *Aurora A*<sup>5</sup>) and alters cofactor binding (eg, *TCF4* binding represses  $\beta$ -catenin pathway genes<sup>6</sup>). Hence, loss of *SMARCB1* creates a malignant potential through activation of different pathways (*SHH*, *Wnt*/ $\beta$ -catenin, *mTOR/Akt*) and loss of cell cycle control.

In *The Lancet Oncology*, Jonathon Torchia and colleagues<sup>7</sup> report the largest integrated molecular and clinicopathological analysis of childhood atypical teratoid rhabdoid tumours. Gene expression and copy number profiling in a discovery cohort of 43 atypical teratoid rhabdoid tumours identified two molecularly distinct subgroups. Group 1 tumours had substantial overexpression of genes involved in the neural development (*FABP7* and *ASCL1*) and *NOTCH* signalling (*HES5/6* and *DLL1/3*) and were predominantly supratentorial. Group 2 atypical teratoid

rhabdoid tumours had enrichment of genes involved in mesenchymal differentiation, cell adhesion and migration, and activation of bone morphogenetic protein (BMP) and MAPK signalling and were generally infratentorial. A validation cohort of 144 primary atypical teratoid rhabdoid tumours showed that robust *ASCL1* antibody immunostaining reliably differentiated the two molecular groups. In a univariate analysis of 70 patients who received treatment with curative intent, the investigators reported that *ASCL1* expression was prognostic of better 5-year overall survival than was no expression (35% [95% CI 13–57] vs 20% [6–34],  $p=0.033$ ). Interestingly, among patients treated with chemotherapy alone, group 1 patients had significantly better 5-year overall survival (34% [7–61]) than did group 2 patients (9% [0–21];  $p=0.001$ ), implying greater intrinsic chemosensitivity in group 1 patients, justifying a radiation-sparing approach in young patients in this group to avoid neurocognitive sequelae, without compromising survival. Extent of resection was significantly greater in group 1 patients irrespective of location, indicating enrichment of cell adhesion and migration genes in group 2 tumours that might promote a more infiltrative or metastatic tumour phenotype.

Integrating molecular and clinicopathological analyses, the investigators proposed three separate risk categories of atypical teratoid rhabdoid tumours: the average risk group comprised older children with completely resected, supratentorial atypical teratoid rhabdoid tumours, high *ASCL1* expression with a significantly better 5-year overall survival of 60% (95% CI 17–100) than in the other categories; very high risk comprised younger children with incompletely resected or metastatic infratentorial, *ASCL1*-negative atypical teratoid rhabdoid tumours with a dismal 5-year overall survival of 6% (0–16) and rapid disease progression; and high risk comprised children with a 5-year overall survival of 32% (95% CI 14–50), which included patients with all other features. Importantly, average-risk patients treated with chemotherapy alone had the best outcomes, whereas highest-risk patients had a dismal prognosis irrespective of therapy.

This study has several limitations. In both the discovery and validation cohorts, full clinical and



Ra Novosti/Science Photo Library

*Lancet Oncol* 2015

Published Online

April 14, 2015

[http://dx.doi.org/10.1016/S1470-2045\(15\)70170-1](http://dx.doi.org/10.1016/S1470-2045(15)70170-1)

S1470-2045(15)70170-1

See Online/Articles

[http://dx.doi.org/10.1016/S1470-2045\(15\)70114-2](http://dx.doi.org/10.1016/S1470-2045(15)70114-2)

S1470-2045(15)70114-2

treatment data were unavailable for many patients. This multinational cohort spanned decades and diverse treatment approaches. Only univariate analyses adjusted for ASCL1 status were used to assess the prognostic significance of molecular subtyping because the cohort was underpowered for multivariate analyses. In risk-stratified analyses, the average-risk cohort was small (n=6). Prospective validation of the proposed risk-stratification in current protocols for atypical teratoid rhabdoid tumours is crucial to confirm these results and refine group-specific diagnostic molecular markers.

Nevertheless, this important study marks the first step towards an integrated, risk-adapted, therapeutic approach for atypical teratoid rhabdoid tumours: avoiding radiation in younger, average-risk patients and integrating novel therapies early for very high-risk patients. Inhibitors of NOTCH, BMP, and LIN28/RAS/MAP kinase signalling are candidates with promising preclinical efficacy in models of atypical teratoid rhabdoid tumours. The MEK inhibitor selumetinib,<sup>8</sup> CDK,<sup>9</sup> Aurora kinase,<sup>10</sup> and EZH2 inhibitors<sup>11</sup> have also shown encouraging specific preclinical activity against atypical teratoid rhabdoid tumours, leading to current or planned paediatric trials. Indeed, efficacy has been reported with the Aurora kinase A inhibitor alisertib in children with atypical teratoid rhabdoid tumours. Rapid validation of these results might herald a promising new therapeutic era of risk-adapted trials incorporating novel agents for this group of children.

Lindsey M Hoffman, \*Maryam Fouladi

Cincinnati Children's Hospital Medical Center, 3333 Burnet Avenue,  
Cincinnati, OH 45229, USA  
maryam.fouladi@cchmc.org

We declare no competing interests.

- 1 Chi SN, Zimmerman MA, Yao X, et al. Intensive multimodality treatment for children with newly diagnosed CNS atypical teratoid rhabdoid tumor. *J Clin Oncol* 2009; **27**: 385–89.
- 2 Lee RS, Roberts CWM. Rhabdoid tumors: an initial clue to the role of chromatin remodeling in cancer. *Brain Pathol* 2013; **23**: 200–05.
- 3 Lee RS, Stewart C, Carter SL, et al. A remarkably simple genome underlies highly malignant pediatric rhabdoid cancers. *J Clin Invest* 2012; **122**: 2983–88.
- 4 Jagani Z, Mora-Blanco EL, Sansam CG, et al. Loss of the tumor suppressor Snf5 leads to aberrant activation of the Hedgehog-Gli pathway. *Nat Med* 2010; **16**: 1429–33.
- 5 Lee S, Cimica V, Ramachandra N, Zagzag D, Kalpana GV. Aurora A is a repressed effector target of the chromatin remodeling protein INI1/hSNF5 required for rhabdoid tumor cell survival. *Cancer Res* 2011; **71**: 3225–35.
- 6 Mora-Blanco EL, Mishina Y, Tillman EJ, et al. Activation of  $\beta$ -catenin/TCF targets following loss of the tumor suppressor SNF5. *Oncogene* 2014; **33**: 933–38.
- 7 Torchia J, Picard D, Lafay-Cousin L, et al. Molecular subgroups of atypical teratoid rhabdoid tumours in children: an integrated genomic and clinicopathological analysis. *Lancet Oncol* 2015; published online April 14. [http://dx.doi.org/10.1016/S1470-2045\(15\)70114-2](http://dx.doi.org/10.1016/S1470-2045(15)70114-2).
- 8 Weingart MF, Roth JJ, Hutt-Cabezas M, et al. Disrupting LIN28 in atypical teratoid rhabdoid tumors reveals the importance of the mitogen activated protein kinase pathway as a therapeutic target. *Oncotarget* 2015; **6**: 3165–77.
- 9 Smith ME, Cimica V, Chinni S, et al. Therapeutically targeting cyclin D1 in primary tumors arising from loss of Ini1. *Proc Natl Acad Sci USA* 2011; **108**: 319–24.
- 10 Wetmore C, Boyett J, Li S, et al. Alisertib is active as single agent in recurrent atypical teratoid rhabdoid tumors in 4 children. *Neuro-Oncology* 2015; published online Feb 16. DOI:10.1093/neuonc/nov017.
- 11 Knutson SK, Warholic NM, Wigle TJ, et al. Durable tumor regression in genetically altered malignant rhabdoid tumors by inhibition of methyltransferase EZH2. *Proc Natl Acad Sci USA* 2013; **110**: 7922–27.